



Esperion to Report Second Quarter 2019 Financial Results August 8, 2019

July 30, 2019

ANN ARBOR, Mich., July 30, 2019 (GLOBE NEWSWIRE) -- Esperion (NASDAQ:ESPR) today announced it will report second quarter 2019 financial results on August 8, 2019 before the U.S. financial markets open.

Bempedoic Acid

Bempedoic acid is our lead, non-statin, orally available, once-daily, LDL-C lowering therapy. With a targeted mechanism of action, bempedoic acid is a first-in-class, ATP Citrate Lyase (ACL) inhibitor that reduces cholesterol biosynthesis and lowers LDL-C by up-regulating the LDL receptor. Similar to statins, bempedoic acid also reduces hsCRP, a key marker of inflammation associated with cardiovascular disease. Completed Phase 3 studies conducted in more than 4,000 patients, with over 2,600 patients treated with bempedoic acid, demonstrated up to 18 percent placebo corrected LDL-C lowering when used with moderate- and high-intensity statins and 21 to 28 percent placebo corrected LDL-C lowering when used with low dose or no background statin.

Bempedoic Acid / Ezetimibe Combination Pill

Through the complementary mechanisms of action of inhibition of cholesterol synthesis (bempedoic acid) and inhibition of cholesterol absorption (ezetimibe), the bempedoic acid / ezetimibe combination tablet is a non-statin, orally available, once-daily, LDL-C lowering therapy. Inhibition of ATP Citrate Lyase (ACL) by bempedoic acid reduces cholesterol biosynthesis and lowers LDL-C by up-regulating the LDL receptor. Inhibition of Niemann-Pick C1-Like 1 (NPC1L1) by ezetimibe results in reduced absorption of cholesterol from the gastrointestinal tract, thereby reducing delivery of cholesterol to the liver, which in turn upregulates the LDL receptors. Phase 3 data demonstrated that this combination results in a 29 percent placebo corrected LDL-C lowering when used with maximally tolerated statins, a 44 percent LDL-C lowering when used with no background statin (post-hoc analysis), and a 34 percent reduction in high sensitivity C-reactive protein (hsCRP).

CLEAR Outcomes

The effect of bempedoic acid on cardiovascular morbidity and mortality has not yet been determined. Esperion initiated a global cardiovascular outcomes trial (CVOT) to assess the effects of bempedoic acid on the occurrence of major cardiovascular events in patients with, or at high risk for, cardiovascular disease (CVD) who are only able to tolerate less than the lowest approved daily starting dose of a statin and considered "statin averse." The CVOT — known as CLEAR Outcomes — is an event-driven, global, randomized, double-blind, placebo-controlled study expected to enroll approximately 12,600 patients with hypercholesterolemia and high CVD risk at over 1,000 sites in approximately 30 countries.

Esperion's Commitment to Patients with Hypercholesterolemia

High levels of LDL-C can lead to a build-up of fat and cholesterol in and on artery walls (known as atherosclerosis), potentially leading to cardiovascular events, including heart attack or stroke. In the U.S., 96 million people, or more than 37 percent of the adult population, have elevated LDL-C. There are approximately 18 million people in the U.S. with atherosclerotic cardiovascular disease (ASCVD) who live with elevated levels of LDL-C despite taking maximally tolerated lipid-modifying therapy — including individuals considered statin averse — leaving them at high risk for cardiovascular events¹. More than 50 percent of ASCVD patients who are not able to reach their LDL-C goals with statins alone need less than a 40 percent reduction to reach their LDL-C threshold².

Esperion's mission as the Lipid Management Company is to deliver once-daily, oral therapies that complement existing oral drugs to provide the additional LDL-C lowering that these patients need.

The Lipid Management Company

Esperion is the Lipid Management Company passionately committed to developing and commercializing convenient, cost-effective, complementary, once-daily, oral therapies for the treatment of patients with elevated LDL-C. Through scientific and clinical excellence, and a deep understanding of cholesterol biology, the experienced Lipid Management Team at Esperion is committed to developing new LDL-C lowering therapies that will make a substantial impact on reducing global cardiovascular disease, the leading cause of death around the world. Bempedoic acid and the bempedoic acid / ezetimibe combination tablet are targeted therapies that are being developed to significantly lower elevated LDL-C levels in patients with hypercholesterolemia, including patients inadequately treated with current lipid-modifying therapies. For more information, please visit www.esperion.com and follow us on Twitter at <https://twitter.com/EsperionInc>.

Forward-Looking Statements

This press release contains forward-looking statements that are made pursuant to the safe harbor provisions of the federal securities laws, including statements regarding the regulatory approval pathway for bempedoic acid and the bempedoic acid / ezetimibe combination tablet and the therapeutic potential of, clinical development plan for, bempedoic acid and the bempedoic acid / ezetimibe combination tablet, including Esperion's timing, designs, plans and announcement of results regarding its CLEAR Outcomes study and other ongoing clinical studies for bempedoic acid and the bempedoic acid / ezetimibe combination tablet, timing for the review and approval of the NDAs and the MAAs and Esperion's expectations for the market for therapies to lower LDL-C, including the market adoption of bempedoic acid and the bempedoic acid / ezetimibe combination tablet, if approved, and the expected upcoming milestones described in this press release. Any express or implied statements contained in this press release that are not statements of historical fact may be deemed to be forward-looking statements. Forward-looking statements involve risks and uncertainties that could cause Esperion's actual results to differ significantly from those projected, including, without limitation, delays or failures in Esperion's studies, that positive results from a clinical study of bempedoic acid may not be sufficient for FDA or EMA approval or necessarily be predictive of the results of future or ongoing clinical studies, that notwithstanding the completion of Esperion's Phase 3 clinical development program for LDL-C

lowering, the FDA or EMA may require additional development in connection with seeking regulatory approval, that existing cash resources may be used more quickly than anticipated, and the risks detailed in Esperion's filings with the Securities and Exchange Commission. Esperion disclaims any obligation or undertaking to update or revise any forward-looking statements contained in this press release, other than to the extent required by law.

References

¹ Esperion market research on file: research project interviewing 350 physicians. Esperion Therapeutics, Inc. Sept-Oct 2018.

² Data on file: analysis of NHANES database. Esperion Therapeutics, Inc. 2018.

Investor Contact:

Alex Schwartz

Esperion

734-249-3386

aschwartz@esperion.com

Media Contact:

Elliot Fox

W2Opure

212-257-6724

efox@purecommunications.com